

Week 5 and 6 Notes

Continuity

If a function is continuous, it must pass these 3 rules

1. c is in the domain of $f(x)$
2. $f(c)$ exists
3. $\lim_{x \rightarrow c} f(x) = f(c)$

Eg. Write a delta-epsilon proof to show that $f(x) = 2x + 1$ is continuous at $x = 5$.

1. Dom of $f(x) = (-\infty, \infty)$

$$5 \in \text{Dom } f(x)$$

$$2. f(5) = 2(5) + 1 \\ = 11$$

$$3. \text{ Prove } \lim_{x \rightarrow 5} 2x + 1 = 11$$

$$\forall \epsilon > 0, \exists \delta > 0 \mid 0 < |x - 5| < \delta \rightarrow |2x + 1 - 11| < \epsilon$$

$$|2x - 10| < \epsilon$$

$$2|x - 5| < \epsilon$$

$$2\delta = \epsilon$$

$$\delta = \frac{\epsilon}{2}$$

$$|2x - 10| < \epsilon$$

$$2|x - 5|$$

$$= 2\delta$$

$$= 2\left(\frac{\epsilon}{2}\right)$$

$$= \epsilon$$

QED


$$\therefore \lim_{x \rightarrow 5} 2x + 1 = 11$$

$\therefore f(x)$ is continuous at $x = 5$

Types of Discontinuities


1. Removable Discontinuity: If $\lim_{x \rightarrow c} f(x)$ exists but isn't equal to $f(c)$

E.g.




2. Jump Discontinuity: If $\lim_{x \rightarrow c^-} f(x)$ and $\lim_{x \rightarrow c^+} f(x)$ exist but are not equal.

E.g.



3. Infinite Discontinuity: If one or both of $\lim_{x \rightarrow c^-} f(x)$ or $\lim_{x \rightarrow c^+} f(x)$ is infinite.

E.g.



Continuity at a point:

If function f is defined on open interval $(c-p, c+p)$, we say that f is continuous at c if $\lim_{x \rightarrow c} f(x) = f(c)$.

The precise definition of continuity at a point is:

If function f is defined on open interval $(c-p, c+p)$, we say that f is continuous at c if for each $\epsilon > 0$, there exists $\delta > 0$ such that if $|x - c| < \delta$ then $|f(x) - f(c)| < \epsilon$.

One Sided Limits

- $f(x)$ is cont from the right at number c if $\lim_{x \rightarrow c^+} f(x) = f(c)$.

$$\forall \epsilon > 0, \exists \delta > 0 \mid |x - c| < \delta \mid \rightarrow |f(x) - f(c)| < \epsilon$$

- $f(x)$ is cont from the left at number c if $\lim_{x \rightarrow c^-} f(x) = f(c)$

$$\forall \epsilon > 0, \exists \delta > 0 \mid |x - c| < \delta \mid \rightarrow |f(x) - f(c)| < \epsilon$$

Continuity on an Interval

- Function f is cont on open interval (a, b) if it is cont at every number in the interval

$$\text{for all } p \in (a, b): \lim_{x \rightarrow p} f(x) = f(p)$$

- Function f is cont on closed interval $[a, b]$ if it is cont on open interval (a, b) , right cont at a and left cont at b .

$$\text{for all } p \in (a, b): \lim_{x \rightarrow p} f(x) = f(p)$$

$$\lim_{x \rightarrow a^+} f(x) = f(a) \text{ and } \lim_{x \rightarrow b^-} f(x) = f(b)$$

3 Important Consequences of Continuity

1. Cont functions on closed interval $[a, b]$ attain their min and max values on that interval.
2. Values of cont functions on closed interval go through every possible value between $f(a)$ and $f(b)$.
3. Cont functions on closed interval are bounded and attain their upper and lower bounds.

- Extreme Value Theorem:

If $f(x)$ is cont on closed interval $[a, b]$, then there exists some values M and m in $[a, b]$ such that $f(M)$ is the max value and $f(m)$ is the min value.

Proof:

Lemma

If f is cont on $[a, b]$, then f is bounded on $[a, b]$.

Proof:

Consider $\{x : x \in [a, b] \text{ and } f \text{ is bounded on } [a, x]\}$

It is easy to see that this set is non-empty and bounded above by b . Thus, we can set $c = \text{lub } \{x : f \text{ is bounded on } [a, x]\}$

To argue $c = b$, suppose $c < b$. Since f is cont at c , it is bounded on $[c - \epsilon, c + \epsilon]$ for some $\epsilon > 0$. This means that f is bounded on $[a, c - \epsilon]$ and on $[c - \epsilon, c + \epsilon]$, which means f is bounded on $[a, c + \epsilon]$.

However, this contradicts our choice of c , so $c = b$. This tells us that f is bounded on $[a, x]$, $x < b$. From the continuity of f , we know it is bounded on $[b - \epsilon, b]$, $b - \epsilon < b$. $\therefore f$ is bounded on $[a, b - \epsilon]$ and $[b - \epsilon, b]$ which means f is bounded on $[a, b]$.

QED

Extreme Value Theorem

By the lemma, f is bounded on $[a, b]$.

Set $M = \text{lub } \{f(x) : x \in [a, b]\}$

We have to show that there exists a c in $[a, b]$ such that $f(c) = M$

To do this, we set $g(x) = \frac{1}{M - f(x)}$

If $f(x) \neq M$, then $g(x)$ is cont on $[a, b]$ and bounded on $[a, b]$.

However, $g(x)$ cannot be bounded on $[a, b]$, which makes $f(x) = M$.

QED

Side Note *

The proof for m is similar

- Intermediate Value Theorem:

If $f(x)$ is cont on $[a, b]$, then for any k strictly between $f(a)$ and $f(b)$, there exists at least one $c \in (a, b)$ such that $f(c) = k$

Proof:

Lemma:

If $f(x)$ is cont on $[a, b]$ and $f(a) < 0 < f(b)$, then $\exists c \in (a, b)$ for which $f(c) = 0$

Proof

Let $f(a) < 0 < f(b)$

Since $f(a) < 0$ and $f(x)$ is cont on $[a, b]$, we know a number E such that $f(x)$ is negative on $[a, E]$. Since $[a, E]$ is bounded above by b , b is an upper bound for $[a, E]$. By the LUB Axiom, this set has a Sup. Let's suppose $\text{Sup} \{E : f \text{ is negative on } [a, E]\} = c$.

We know that $c \leq b$, but since $f(x)$ is negative on $[a, E]$ and $f(b) > 0$, $c \neq b$. Since $c < b$, $f(c) < 0$ or $f(c) = 0$. However, if $f(c) < 0$, then there exists a number $c' \in E$ for which $f(c') < 0$ and $c' \in$ becomes the new Sup. $\therefore f(c) = 0$

QED

Proof of IVT:

Let $g(x) = f(x) - k$ for any $f(a) < k < f(b)$

Then, $g(a) = f(a) - k < 0$ and $g(b) = f(b) - k > 0$

From the lemma, we know $\exists x = c$, such that $g(c) = 0$.


$$\therefore g(c) = f(c) - k$$

$$0 = f(c) - k$$


$$f(c) = k$$

QED

- Extreme Value Theorem: If f is continuous on a closed interval $[a, b]$, then there exists values M and m in the interval $[a, b]$ such that $f(m)$ is the max value and $f(m)$ is the min value.



- Intermediate Value Theorem: If f is cont. on a closed interval $[a, b]$, then for any k between $f(a)$ and $f(b)$, there exists a $c \in (a, b)$ such that $f(c) = k$.



- A function can only change signs at a point $x=c$ only if $f(x)=0$, undefined or discontinuous at $x=c$.

- Least Upper Bound Axiom / Sup:

Every nonempty set of real numbers that has an upper bound has at a LUB. The LUB may not directly be in the set, but can be found from the pattern.

E.g. $S = \left\{ \frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \frac{5}{6}, \dots \right\}$

The LUB of S is 1, because the numbers are approaching 1.

The largest element of a set, $\max S$, exists if LUB S exists and LUB $S \in S$.

- If M is the LUB of set S and ϵ is a positive number, then there exists at least 1 number s in S such that $M - \epsilon < s \leq M$.

Proof: Let $\epsilon > 0$

$$M - \epsilon < s$$

Suppose on the contrary there is no number in S .

That means $x \leq M - \epsilon$ for all $x \in S$.

This makes $M - \epsilon$ an LUB that is smaller than M , which can't be true.

$\therefore$ Our supposition is wrong.

QED

- Every nonempty set of real numbers that has a lower bound has a GLB or Inf.

Proof: Suppose S is nonempty and has a lower bound k .

$$k \leq s, \text{ for all } s \in S$$

$$-s \leq -k, \text{ for all } s \in S$$

From the LUB axiom, we conclude that

$\{-s : s \in S\}$ has a LUB, m .

$$-s \leq m \text{ for all } s \in S$$

$$-m \leq s \text{ for all } s \in S$$

Suppose $-m < m_1 \leq s$, for all $s \in S$.

$$-s \leq -m_1 < m \text{ for all } s \in S$$

Thus m would not be LUB of $\{-s : s \in S\}$

- If m is the GLB of set S and ϵ is a positive number, then there exists a number s in S such that $m \leq s < m + \epsilon$

(3)

- Bound Value Theorem:

If f is cont. on $[a, b]$, then f is bounded on $[a, b]$.

Proof: Consider $\{x: x \in [a, b] \text{ and } f \text{ is bounded on } [a, x]\}$

Set $c = \text{LUB } \{x: f \text{ is bounded on } [a, x]\}$

Suppose $c < b$. From the continuity of f at c , f is bounded on $[c - \epsilon, c + \epsilon]$, $\epsilon > 0$. But this contradicts our choice of c . So $c = b$. This tells us

f is bounded on $[a, x]$ with $x < b$. From the continuity of f , we know that f is bounded on some interval of the form $[b - \epsilon, b]$. Since $b - \epsilon < b$, f is bounded on $[a, b - \epsilon]$ and $[b - \epsilon, b]$. $\therefore f$ is bounded on $[a, b]$

- Properties of Limits

1. The sum/difference of 2 cont. functions is cont.
2. The product/quotient of 2 cont. functions is cont.
3. The composition (fog) of 2 cont functions is cont.
4. All polynomial functions are cont.

Limit Proofs

Let $\lim_{x \rightarrow c} f(x) = L$ and $\lim_{x \rightarrow c} g(x) = M$

1. Constant Multiple Law

Prove $\lim_{x \rightarrow c} k f(x) = k \lim_{x \rightarrow c} f(x)$

$$\lim_{x \rightarrow c} k f(x)$$

$$= \lim_{x \rightarrow c} k \cdot \lim_{x \rightarrow c} f(x)$$

$$= k \lim_{x \rightarrow c} f(x)$$

2. Addition Law

Prove $\lim_{x \rightarrow c} [f(x) + g(x)] = \lim_{x \rightarrow c} f(x) + \lim_{x \rightarrow c} g(x)$

$\lim_{x \rightarrow c} f(x) = L$ means $\forall \epsilon_1 > 0, \exists d_1 > 0 \mid 0 < |x - c| < d_1 \Rightarrow |f(x) - L| < \epsilon_1$

Let ϵ_1 be $\frac{\epsilon}{2}$

$\lim_{x \rightarrow c} g(x) = M$ means $\forall \epsilon_2 > 0, \exists d_2 > 0 \mid 0 < |x - c| < d_2 \Rightarrow |g(x) - M| < \epsilon_2$

Let ϵ_2 be $\frac{\epsilon}{2}$

Choose d to be $\min(d_1, d_2)$

Then, both $f(x)$ and $g(x)$ are within $\frac{\epsilon}{2}$ of L and M .

$$-\frac{\epsilon}{2} < f(x) - L < \frac{\epsilon}{2}$$

$$+ -\frac{\epsilon}{2} < g(x) - M < \frac{\epsilon}{2}$$

$$- \epsilon < f(x) + g(x) - (L + M) < \epsilon$$

QED

3. Product Rule * Optional

Prove $\lim_{x \rightarrow c} [f(x) \cdot g(x)] = LM$

$\lim_{x \rightarrow c} [f(x) - L] = 0$ and $\lim_{x \rightarrow c} [g(x) - M] = 0$

Proof: $\lim_{x \rightarrow c} [f(x) - L]$

$$= \lim_{x \rightarrow c} f(x) - \lim_{x \rightarrow c} L$$

$$= L - L$$

$$= 0$$

The same logic applies to $\lim_{x \rightarrow c} [g(x) - M] = 0$.

(4)

Now, we have to prove $\lim_{x \rightarrow c} [f(x) - L][g(x) - M] = 0$
 $\lim_{x \rightarrow c} [f(x) - L] = 0$ means:

$\forall \epsilon_1 > 0, \exists \delta_1 > 0 \mid 0 < |x - c| < \delta_1 \rightarrow |f(x) - L| < \epsilon_1$

$\lim_{x \rightarrow c} [g(x) - M] = 0$ means:

$\forall \epsilon_2 > 0, \exists \delta_2 > 0 \mid 0 < |x - c| < \delta_2 \rightarrow |g(x) - M| < \epsilon_2$

Let $\epsilon_1 = \sqrt{\epsilon}$ and $\epsilon_2 = \sqrt{\epsilon}$

Choose δ to be $\min(\delta_1, \delta_2)$

Then: $|f(x) - L| < \sqrt{\epsilon}$ and $|g(x) - M| < \sqrt{\epsilon}$

Multiplying the two together, we get $|f(x) - L| |g(x) - M| < \epsilon$

$\therefore \lim_{x \rightarrow c} [f(x) - L][g(x) - M] = 0$

If we multiply $[f(x) - L][g(x) - M]$, we get $f(x)g(x) - f(x)M - g(x)L + LM$

$$f(x)g(x) = [f(x) - L][g(x) - M] + f(x)M + g(x)L - LM$$

$$\lim_{x \rightarrow c} f(x)g(x) = \lim_{x \rightarrow c} [f(x) - L][g(x) - M] + \lim_{x \rightarrow c} f(x)M + \lim_{x \rightarrow c} g(x)L - \lim_{x \rightarrow c} LM$$

$$= \lim_{x \rightarrow c} [f(x) - L][g(x) - M] + \lim_{x \rightarrow c} f(x)M + \lim_{x \rightarrow c} g(x)L - \lim_{x \rightarrow c} LM$$

$$= 0 + LM + LM - LM$$

$$= LM$$

QED

4. Difference Law

Prove $\lim_{x \rightarrow c} [f(x) - g(x)] = L - M$

$$\begin{aligned} & \lim_{x \rightarrow c} [f(x) - g(x)] \\ &= \lim_{x \rightarrow c} [f(x) + (-g(x))] \\ &= \lim_{x \rightarrow c} f(x) + \lim_{x \rightarrow c} (-g(x)) \\ &= \lim_{x \rightarrow c} f(x) - \lim_{x \rightarrow c} g(x) \\ &= L - M \end{aligned}$$

QED

5. Reciprocal Law

Prove $\lim_{x \rightarrow c} \frac{1}{g(x)} = \frac{1}{M}, M \neq 0$

$$\forall \epsilon > 0, \exists \delta > 0 \text{ s.t. } 0 < |x - c| < \delta \Rightarrow \left| \frac{1}{g(x)} - \frac{1}{M} \right| < \epsilon$$

$$\begin{aligned} & \left| \frac{1}{g(x)} - \frac{1}{M} \right| \\ &= \left| \frac{M - g(x)}{M(g(x))} \right| \quad (|M - g(x)| = |g(x) - M| \text{ Side Note}) \\ &= \left| \frac{1}{M} \right| \left| \frac{1}{g(x)} \right| |g(x) - M| \end{aligned}$$

$$\text{Choose } d_1 \text{ such that } |g(x) - M| < \frac{|M|}{2} \Rightarrow |g(x)| > \frac{|M|}{2} \Rightarrow \frac{1}{|g(x)|} < \frac{2}{|M|}$$

$$\text{Choose } d_2 \text{ such that } |g(x) - M| < \frac{M^2 \epsilon}{2}$$

Choose d to be $\min(d_1, d_2)$

$$\begin{aligned} \left| \frac{1}{M} \right| \left| \frac{1}{g(x)} \right| |g(x) - M| &< \frac{2}{M^2} \left(\frac{M^2}{2} \right) (\epsilon) \\ &< \epsilon \end{aligned}$$

QED

6. Quotient Law * Optional

(5)

Prove $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{L}{M}, M \neq 0$

We can rewrite $\lim_{x \rightarrow c} \frac{f(x)}{g(x)}$ as $\lim_{x \rightarrow c} f(x) \left(\frac{1}{g(x)} \right)$

$$\lim_{x \rightarrow c} f(x) \left(\frac{1}{g(x)} \right)$$

$$= \lim_{x \rightarrow c} f(x) \cdot \lim_{x \rightarrow c} \frac{1}{g(x)} \quad \leftarrow \text{We proved this in the reciprocal law}$$

$$= L \cdot \frac{1}{M}$$

$$= \frac{L}{M} \quad \text{QED}$$

7. Continuity of power function

Prove that $f(x) = x^k$ is continuous on $(-\infty, \infty)$

We have to prove $f(x)$ is cont on $(-\infty, \infty)$ first

1. $\text{Dom } f(x) = (-\infty, \infty)$

2. $f(c) = c^k$

3. $\lim_{x \rightarrow c} x^k = f(c) = c^k$

By the product rule, $\lim_{x \rightarrow c} x^k = \lim_{x \rightarrow c} x \cdot \lim_{x \rightarrow c} x \cdots$

$$= c \cdot c \cdot c \cdots$$

$$= c^k$$

QED

* Side Note *

We can also prove the continuity of $f(x) = x^{-k}$ and $f(x) = x^{p/q}$ using the same strategy.

9. Squeeze Theorem

6

Let $p > 0$. Suppose that $\forall x$ such that $0 < |x - c| < p$, $h(x) \leq f(x) \leq g(x)$.
If $\lim_{x \rightarrow c} h(x) = L$ and $\lim_{x \rightarrow c} g(x) = L$, then $\lim_{x \rightarrow c} f(x) = L$

Proof: $\forall \epsilon > 0, \exists d_1 > 0 \mid 0 < |x - c| < d_1 \rightarrow |h(x) - L| < \epsilon$

$\forall \epsilon > 0, \exists d_2 > 0 \mid 0 < |x - c| < d_2 \rightarrow |g(x) - L| < \epsilon$

Choose d to be $\min(d_1, d_2)$

Then, $|h(x) - L| < \epsilon$ and $|g(x) - L| < \epsilon$

↓

↓

$$-\epsilon < h(x) - L < \epsilon$$

$$-\epsilon < g(x) - L < \epsilon$$

$$L - \epsilon < h(x) < L + \epsilon$$

$$L - \epsilon < g(x) < L + \epsilon$$

Since $L - \epsilon < h(x) \leq f(x) \leq g(x) < L + \epsilon$,

then $L - \epsilon < f(x) < L + \epsilon$


QED

10. Prove the continuity of $f(x) = \sin(x)$

1. Domain $\sin(x) = (-\infty, \infty)$

2. $f(c) = \sin(c)$

3. Find $\lim_{x \rightarrow c} \sin x = \sin c$



$$\sin \theta = |AB|$$

$$0 \leq |AB| \leq \theta$$

$$\lim_{\theta \rightarrow 0} 0 = 0 \quad \text{and} \quad \lim_{\theta \rightarrow 0} \theta = 0$$

$$\therefore \text{By S.T., } \lim_{\theta \rightarrow 0} \sin \theta = 0$$

We also have to prove $\lim_{\theta \rightarrow 0} \cos \theta = 1$

$$0 \leq |BC| \leq \theta$$

$$0 \leq |1 - \cos \theta| \leq \theta$$

$$-1 \leq -\cos \theta \leq \theta - 1$$

$$1 \geq \cos \theta \geq 1 - \theta$$

$$\lim_{\theta \rightarrow 0} 1 = 1 \quad \text{and} \quad \lim_{\theta \rightarrow 0} 1 - \theta = 1$$

$$\therefore \text{By S.T., } \lim_{\theta \rightarrow 0} \cos \theta = 1$$

Now, sub $x = cth$ in. If $x \rightarrow c$, then $h \rightarrow 0$

$$\lim_{x \rightarrow c} \sin x$$

$$= \lim_{h \rightarrow 0} \sin(cth)$$

$$= \lim_{h \rightarrow 0} (\sin c \cosh + \sinh \cos c)$$

$$= \lim_{h \rightarrow 0} \sin c \cdot \cosh + \lim_{h \rightarrow 0} \sinh \cos c$$

$$= \lim_{h \rightarrow 0} \sin c \cdot \lim_{h \rightarrow 0} \cosh + \lim_{h \rightarrow 0} \sinh \cdot \lim_{h \rightarrow 0} \cos c$$

$$= \sin c$$

QED

11. Prove the continuity of $f(x) = e^x$

We define e to be the number that $(1+h)^{\frac{1}{h}}$ approaches as h approaches 0.

$$\lim_{h \rightarrow 0} (1+h)^{\frac{1}{h}} = e \quad \text{and} \quad \lim_{h \rightarrow 0} \left(\frac{e^h - 1}{h} \right) = 1$$

$$1. \text{ Dom } e^x = (-\infty, \infty)$$

$$2. f(c) = e^c$$

$$3. \lim_{x \rightarrow c} e^x = e^c$$

Sub $x = cth$

$$\lim_{h \rightarrow 0} e^x = \lim_{h \rightarrow 0} e^{cth}$$

$$= \lim_{h \rightarrow 0} e^c \cdot \lim_{h \rightarrow 0} e^h$$

$$= e^c \cdot \lim_{h \rightarrow 0} (e^h + 1 - 1)$$


$$= e^c \cdot \lim_{h \rightarrow 0} \left(\frac{e^h - 1}{h} h + 1 \right)$$

$$\rightarrow = e^c \left(\lim_{h \rightarrow 0} \frac{e^h - 1}{h} \cdot \lim_{h \rightarrow 0} h + \lim_{h \rightarrow 0} 1 \right)$$

$$= e^c$$

QED

12 Prove $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$



$$\text{Area } \triangle OPA = \frac{\sin \theta}{2}$$

$$\text{Area Sector OPA} = \frac{\theta}{2}$$

Area of $\triangle OPA \leq \text{Area of Sector OPA} \leq \text{Area OQA}$

$$\frac{\sin \theta}{2} \leq \frac{\theta}{2} \leq \frac{\tan \theta}{2}$$

$$\sin \theta \leq \theta \leq \tan \theta$$

$$1 \leq \frac{\theta}{\sin \theta} \leq \frac{1}{\cos \theta}$$

$$1 \geq \frac{\sin \theta}{\theta} \geq \cos \theta$$

$$\lim_{x \rightarrow 0} 1 = 1 \quad \text{and} \quad \lim_{x \rightarrow 0} \theta = 1$$

$$\therefore \text{By S.T., } \lim_{x \rightarrow 0} \frac{\sin \theta}{\theta} = 1$$

QED

13 Prove $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x} = 0$

$$\begin{aligned} & \lim_{x \rightarrow 0} \frac{1 - \cos x}{x} \left(\frac{1 + \cos x}{1 + \cos x} \right) \\ &= \lim_{x \rightarrow 0} \frac{\sin^2 x}{x(1 + \cos x)} \\ &= \lim_{x \rightarrow 0} \frac{\sin x}{x} \cdot \lim_{x \rightarrow 0} \frac{\sin x}{1 + \cos x} \\ &= 0 \\ &\text{QED} \end{aligned}$$

14 Prove if $\lim_{x \rightarrow c} \frac{f(x)}{g(x)}$ is of the form $\frac{1}{0^+}$, then $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \infty$

$\lim_{x \rightarrow c} f(x) = 1$ means

$$\begin{aligned} \forall \epsilon_1 > 0, \exists d_1 > 0 \mid 0 < |x - c| < d_1 \Rightarrow |f(x) - 1| < \epsilon_1 \\ \downarrow \\ 1 - \epsilon_1 < f(x) < 1 + \epsilon_1 \end{aligned}$$

$\lim_{x \rightarrow c} g(x) = 0$ means

$$\begin{aligned} \forall \epsilon_2 > 0, \exists d_2 > 0 \mid 0 < |x - c| < d_2 \Rightarrow |g(x)| < \epsilon_2 \\ \downarrow \\ -\epsilon_2 < g(x) < \epsilon_2 \end{aligned}$$

Choose d to be $\min(d_1, d_2)$.

That means $\frac{f(x)}{g(x)} > \frac{1 - \epsilon_1}{\epsilon_2}$

Let ϵ_1 be $\frac{1}{2}$

Let ϵ_2 be $\frac{1}{2M}$

$$\frac{f(x)}{g(x)} > \frac{1/2}{1/2M}$$

$> M$

QED

15. If $\lim_{x \rightarrow c} \frac{f(x)}{g(x)}$ is in the form of $\frac{1}{0^-}$, then $\frac{f(x)}{g(x)} = -\infty$

$\lim_{x \rightarrow c} f(x) = 1$ means

$$\forall \epsilon_1 > 0, \exists \delta_1 > 0 \mid 0 < |x - c| < \delta_1 \Rightarrow 1 - \epsilon_1 < f(x) < 1 + \epsilon_1$$

$\lim_{x \rightarrow c} g(x) = 0$ means

$$\forall \epsilon_2 > 0, \exists \delta_2 > 0 \mid 0 < |x - c| < \delta_2 \Rightarrow -\epsilon_2 < g(x) < \epsilon_2$$

Choose δ to be $\min(\delta_1, \delta_2)$.

$$\text{Then } \frac{f(x)}{g(x)} > \frac{1 - \epsilon_1}{\epsilon_2}$$

Let ϵ_1 be $\frac{1}{2}$.

Let ϵ_2 be $\frac{1}{2M}$

$$\frac{f(x)}{g(x)} > M$$

QED

16. If $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$ is of the form $\frac{1}{\infty}$, then $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 0$

$\lim_{x \rightarrow \infty} f(x) = 1$ means:

$$\forall \epsilon_1 > 0, \exists N_1 > 0 \mid x > N_1 \Rightarrow 1 - \epsilon_1 < f(x) < 1 + \epsilon_1$$

$\lim_{x \rightarrow \infty} g(x) = \infty$ means:

$$\forall M > 0, \exists N_2 > 0 \mid x > N_2 \Rightarrow g(x) > M$$

Choose N to be $\max(N_1, N_2)$

$$\text{That means } \left| \frac{f(x)}{g(x)} \right| < \frac{1 + \epsilon_1}{M}$$

Let $\epsilon_1 = 1$ and let $M = \frac{2}{\epsilon}$

$$\left| \frac{f(x)}{g(x)} \right| < \epsilon$$

QED

17 If $\lim_{x \rightarrow -\infty} \frac{f(x)}{g(x)}$ is in the form of $\frac{1}{-\infty}$, then $\lim_{x \rightarrow -\infty} \frac{f(x)}{g(x)} = 0$

$\lim_{x \rightarrow -\infty} f(x) = 1$ means

$$\forall \epsilon_1 > 0, \exists N_1 < 0 \mid x < -N_1 \rightarrow 1 - \epsilon_1 < f(x) < 1 + \epsilon_1$$

$\lim_{x \rightarrow -\infty} g(x) = -\infty$ means

$$\forall M < 0, \exists N_2 < 0 \mid x < -N_2 \rightarrow g(x) < -M$$

Choose N to be $\max(N_1, N_2)$

$$\left| \frac{f(x)}{g(x)} \right| < \frac{1 + \epsilon_1}{M}$$

Let $\epsilon_1 = 1$ and let $\epsilon_2 = -2/\epsilon$

$$\left| \frac{f(x)}{g(x)} \right| = M$$

QED

18. Continuity of Composition Functions

If g is cont at c and f is cont at $g(c)$, then $f \circ g$ is cont. at c .
WTS: $\forall \epsilon > 0, \exists \delta > 0 \mid 0 < |x - c| < \delta \rightarrow |f(g(x)) - f(g(c))| < \epsilon$

To show g is cont at c :

$$\exists \delta_1 > 0 \mid 0 < |x - c| < \delta_1 \rightarrow |g(x) - g(c)| < \delta_1 \quad (1)$$

To show f is cont at $g(c)$:

$$\exists \delta_2 > 0 \mid 0 < |g(x) - g(c)| < \delta_2 \rightarrow |f(g(x)) - f(g(c))| < \epsilon \quad (2)$$

Combining 1 and 2, we get

$$\forall \epsilon > 0, \exists \delta > 0 \mid 0 < |x - c| < \delta \rightarrow |f(g(x)) - f(g(c))| < \epsilon$$

QED