

Week 9 Notes

Proof of Trig Derivatives

1. $(\sin x)' = \cos x$

$$\begin{aligned}
 (\sin x)' &= \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sin x \cosh + \sinh \cos x - \sin x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sin x \cosh - \sin x}{h} + \lim_{h \rightarrow 0} \frac{\sinh \cos x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sin x (\cosh - 1)}{h} + \lim_{h \rightarrow 0} \cos x \left(\frac{\sinh}{h} \right) \\
 &= \cos x
 \end{aligned}$$

2. $(\cos x)' = -\sin x$

$$\begin{aligned}
 (\cos x)' &= \lim_{h \rightarrow 0} \frac{\cos(x+h) - \cos x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\cos x \cosh - \sin x \sinh - \cos x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\cos x \cosh - \cos x}{h} - \lim_{h \rightarrow 0} \frac{\sin x \sinh}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\cos x (\cosh - 1)}{h} - \lim_{h \rightarrow 0} \sin x \\
 &= -\sin x
 \end{aligned}$$

3. $(\tan x)' = \sec^2 x$

$$\begin{aligned}
 (\tan x)' &= \lim_{h \rightarrow 0} \frac{\tan(x+h) - \tan x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{\sin(x+h)}{\cos(x+h)} - \frac{\sin x}{\cos x}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sin(x+h) \cos x - \sin x (\cos(x+h))}{h (\cos x) (\cos(x+h))} \\
 &= \lim_{h \rightarrow 0} \frac{\cos x (\sin x \cosh + \cos x \sinh) - \sin x (\cos x \cosh - \sin x \sinh)}{h (\cos x) (\cos(x+h))} \\
 &= \lim_{h \rightarrow 0} \frac{\cos^2 x \sinh + \sin^2 x \sinh}{h (\cos x) (\cos(x+h))}
 \end{aligned}$$

$$= \lim_{h \rightarrow 0} \frac{\sinh(\cos^2 x + \sin^2 x)}{\cos x (\cos(x+h)) h}$$

$$= \frac{1}{\cos^2 x}$$

$$= \sec^2 x$$

$$4. (\sec x)' = \sec x \tan x$$

$$(\sec x)' = \left(\frac{1}{\cos x} \right)'$$

$$= \frac{-(-\sin x)}{\cos^2 x}$$

$$= \frac{\sin x}{\cos^2 x}$$

$$= \sec x \cdot \tan x$$

$$5. (\csc x)' = -\csc x \cdot \cot x$$

$$(\csc x)' = \left(\frac{1}{\sin x} \right)'$$

$$= \frac{-\cos x}{\sin^2 x}$$

$$= -\csc x \cdot \cot x$$

$$6. (\cot x)' = -\csc^2(x)$$

$$(\cot x)' = \left(\frac{1}{\tan x} \right)'$$

$$= \frac{-\sec^2 x}{\tan^2 x}$$

$$= \frac{-1}{\cos^2 x}$$

$$= -\csc^2 x$$

Proof of Inverse Trig Functions

1. $(\arcsin x)' = \frac{1}{\sqrt{1-x^2}}$

$$y = \arcsin(x)$$

$$x = \sin y$$

$$x' = (\sin y)'$$

$$1 = \cos y \cdot y'$$

$$y' = \frac{1}{\cos y}$$

$$= \frac{1}{\sqrt{1-\sin^2 y}} \leftarrow \cos x = \sqrt{1-\sin^2 x}$$

$$= \frac{1}{\sqrt{1-x^2}}$$

2. $(\arccos x)' = \frac{-1}{\sqrt{1-x^2}}$

$$y = \arccos x$$

$$x = \cos y$$

$$x' = (\cos y)'$$

$$1 = -\sin(y) \cdot y'$$

$$y' = \frac{-1}{\sin y}$$

$$= \frac{-1}{\sqrt{1-x^2}}$$

3. $(\arctan x)' = \frac{1}{1+x^2}$

$$y = \arctan x$$

$$x = \tan y$$

$$x' = (\tan y)'$$

$$1 = \sec^2 y \cdot y'$$

$$y' = \frac{1}{\sec^2 y}$$

$$= \frac{1}{1+\tan^2 y} = \frac{1}{1+x^2}$$

$$4. (\operatorname{arc cot} x)' = \frac{-1}{1+x^2}$$

$$y = \operatorname{arc cot} x$$

$$x = \cot y$$

$$x' = (\cot y)'$$

$$1 = -\csc^2 y \cdot y'$$

$$y' = \frac{-1}{\csc^2 y}$$

$$= \frac{-1}{1 + \cot^2 y}$$

$$= \frac{-1}{1+x^2}$$

$$5. (\operatorname{arc sec} x)' = \frac{1}{|x|\sqrt{x^2-1}}$$

$$y = \operatorname{arc sec} x$$

$$x = \sec y$$

$$x' = (\sec y)'$$

$$1 = \sec y \cdot \tan y \cdot y'$$

$$y' = \frac{1}{\sec y \cdot \tan y}$$

$$= \frac{1}{|x|\sqrt{x^2-1}}$$

$$6. (\operatorname{arc csc} x)' = \frac{-1}{|x|\sqrt{x^2-1}}$$

$$y = \operatorname{arc csc} x$$

$$x = \csc y$$

$$x' = (\csc y)'$$

$$1 = -\csc y \cdot \cot y \cdot y'$$



$$y' = \frac{-1}{\csc y \cdot \cot y}$$


$$= \frac{-1}{|x|\sqrt{x^2-1}}$$

Hyperbolic Identities and Proofs

1. $\sinh(t) = \frac{e^t - e^{-t}}{2}$
2. $\cosh(t) = \frac{e^t + e^{-t}}{2}$
3. $\tanh(t) = \frac{\sinh(t)}{\cosh(t)}$
 $= \frac{e^t - e^{-t}}{e^t + e^{-t}}$
4. $\cosh^2 t - \sinh^2 t = 1$

Graph of hyperbolic functions

 $y = \sinh(x)$  $y = \cosh(x)$ 



Prove

1. $(\sinh x)' = \cosh x$

$$\left[\left(\frac{1}{2} \right) (e^x - e^{-x}) \right]'$$

$$= \frac{1}{2} (e^x + e^{-x})$$

$$= \cosh x$$

QED

2. $(\cosh x)' = \sinh x$

$$\left[\frac{1}{2} (e^x + e^{-x}) \right]'$$

$$= \frac{1}{2} (e^x - e^{-x})$$

$$= \sinh x$$

QED

3. $(\tanh x)' = \operatorname{sech}^2 x$

$$\left(\frac{e^x - e^{-x}}{e^x + e^{-x}} \right)'$$

$$= \frac{(e^x + e^{-x})(e^x + e^{-x}) - (e^x - e^{-x})(e^x - e^{-x})}{(e^x + e^{-x})^2}$$

$$= \frac{\cosh^2 x - \sinh^2 x}{\cosh^2 x}$$

$$= \frac{1}{\cosh^2 x}$$

$$= \operatorname{sech}^2 x$$

(4)

$$\begin{aligned}
 4. \quad (\operatorname{sech} x)' &= -\operatorname{sech} x \cdot \tanh x \\
 (\operatorname{sech} x)' &= \left(\frac{1}{\cosh x} \right)' \\
 &= \left(\frac{-\sinh x}{\cosh^2 x} \right) \\
 &= -\operatorname{sech} x \cdot \tanh x \\
 &\quad \text{QED}
 \end{aligned}$$

$$\begin{aligned}
 5. \quad (\operatorname{csch} x)' &= -\operatorname{csch} x \cdot \coth x \\
 (\operatorname{csch} x)' &= \left(\frac{1}{\sinh x} \right)' \\
 &= \frac{-\cosh x}{\sinh^2 x} \\
 &= -\operatorname{csch} x \cdot \coth x \\
 &\quad \text{QED}
 \end{aligned}$$

$$\begin{aligned}
 6. \quad (\coth x)' &= -\operatorname{csch}^2 x \\
 (\coth x)' &= \left(\frac{1}{\tanh x} \right)' \\
 &= \frac{-\operatorname{sech}^2 x}{\tanh^2 x} \\
 &= -\operatorname{csch}^2 x \\
 &\quad \text{QED}
 \end{aligned}$$

$$7. \quad (\sinh^{-1} x)' = \frac{1}{\sqrt{x^2 + 1}}$$

$$\begin{aligned}
 y &= \sinh^{-1} x \\
 x &= \sinh(y) \\
 x' &= \cosh(y) \cdot y' \\
 1 &= \cosh y \cdot y' \\
 y' &= \frac{1}{\cosh y}
 \end{aligned}$$

$$\begin{aligned}
 y' &= \frac{1}{\sqrt{1 + \sinh^2(\sinh^{-1} x)}} \\
 &= \frac{1}{\sqrt{x^2 + 1}} \\
 &\quad \text{QED}
 \end{aligned}$$

$$8. (\cosh^{-1} x)' = \frac{1}{\sqrt{x^2-1}}$$

$$y = \cosh^{-1} x$$

$$x = \cosh y$$

$$x' = (\cosh y)'$$

$$1 = \sinh y \cdot y'$$

$$y' = \frac{1}{\sinh y}$$

$$= \frac{1}{\sqrt{\sinh^2(\cosh^{-1} x) - 1}}$$

$$= \frac{1}{\sqrt{x^2-1}}$$

$$9. (\tanh^{-1} x)' = \frac{1}{1-x^2}$$

$$y = \tanh^{-1} x$$

$$x = \tanh y$$

$$x' = (\tanh y)'$$

$$1 = \operatorname{sech}^2 y \cdot y'$$

$$y' = \frac{1}{\operatorname{sech}^2 y}$$

$$= \frac{1}{\operatorname{sech}^2(\tanh^{-1} x)}$$

$$= \frac{1}{1-x^2}$$

$$10. (\operatorname{sech}^{-1} x)' = \frac{-1}{x\sqrt{1-x^2}}$$

$$y = \operatorname{sech}^{-1} x$$

$$x = \operatorname{sech} y$$

$$x' = (\operatorname{sech} y)'$$

$$1 = -\operatorname{sech} y \cdot \tanh y \cdot y'$$

$$y' = \frac{-1}{\operatorname{sech} y \cdot \tanh y}$$

$$= \frac{-1}{\operatorname{sech}(\operatorname{sech}^{-1} x) \cdot \tanh(\operatorname{sech}^{-1} x)}$$

$$\Rightarrow y' = \frac{-1}{x\sqrt{1-x^2}}$$

$$11. (\operatorname{csch}^{-1} x)' = \frac{-1}{|x| \sqrt{1+x^2}}$$

$$y = \operatorname{csch}^{-1} x$$

$$x = \operatorname{csch} y$$

$$x' = (\operatorname{csch} y)'$$

$$1 = -\operatorname{csch} y \cdot \operatorname{coth} y \cdot y'$$

$$y' = \frac{-1}{\operatorname{csch} y \cdot \operatorname{coth} y}$$

$$= \frac{-1}{\operatorname{csch}(\operatorname{csch}^{-1} x) \cdot \operatorname{coth}(\operatorname{csch}^{-1} x)}$$

$$= \frac{-1}{|x| \sqrt{1+x^2}}$$

$$12. (\operatorname{coth}^{-1} x)' = \frac{1}{1-x^2}$$

$$y = \operatorname{coth}^{-1} x$$

$$x = \operatorname{coth} y$$

$$x' = (\operatorname{coth} y)'$$

$$1 = -\operatorname{csch}^2 y \cdot y'$$

$$y' = \frac{-1}{\operatorname{csch}^2 y}$$

$$= \frac{-1}{\operatorname{csch}^2(\operatorname{coth}^{-1} x)}$$

$$= \frac{-1}{x^2 - 1} = \frac{1}{1-x^2}$$

$$13. \sinh^{-1} x = \ln(x + \sqrt{x^2 + 1}) \text{ for any } x$$

$$\text{let } y = \sinh^{-1} x$$

$$x = \sinh y$$

$$= \frac{e^y - e^{-y}}{2}$$

$$2x = e^y - e^{-y}$$

$$= e^{\sinh^{-1} x} - e^{-\sinh^{-1} x}$$

$$2x e^{\sinh^{-1} x} = (e^{\sinh^{-1} x} - e^{-\sinh^{-1} x}) e^{\sinh^{-1} x}$$

$$= (e^{\sinh^{-1} x})^2 - 1$$

Let u be $e^{\sinh^{-1} x}$

$$0 = u^2 - 2xu - 1$$

$$u = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{2x \pm \sqrt{4x^2 + 4}}{2}$$

$$= x \pm \sqrt{x^2 + 1}$$

However, since $x - \sqrt{x^2 + 1} < 0$, we reject this case

$$\therefore u = x + \sqrt{x^2 + 1}$$

$$e^{\sinh^{-1} x} = x + \sqrt{x^2 + 1}$$

$$\ln(u) = \ln(x + \sqrt{x^2 + 1})$$

$$\sinh^{-1} x = \ln(x + \sqrt{x^2 + 1})$$

14. $\cosh^{-1} x = \ln(x + \sqrt{x^2 - 1})$ for $x \geq 1$ (Proof is similar to 13)

Let $y = \cosh^{-1} x$

$$x = \cosh y$$

$$= \frac{e^y + e^{-y}}{2}$$

$$2x = e^{\cosh^{-1} x} + e^{-\cosh^{-1} x}$$

$$2xe^{\cosh^{-1} x} = (e^{\cosh^{-1} x} + e^{-\cosh^{-1} x})e^{\cosh^{-1} x}$$

$$= (e^{\cosh^{-1} x})^2 + 1$$

$$\text{Let } u = e^{\cosh^{-1} x}$$

$$0 = u^2 - 2xu + 1$$

$$u = \frac{2x \pm \sqrt{4x^2 - 4}}{2}$$

$$= x \pm \sqrt{x^2 - 1}$$

Since $x - \sqrt{x^2 - 1} < 0$, we reject this case

$$u = x + \sqrt{x^2 - 1}$$

$$\ln(u) = \ln(x + \sqrt{x^2 - 1})$$

$$\cosh^{-1} x = \ln(x + \sqrt{x^2 - 1})$$

15 $\tanh^{-1} x = \frac{1}{2} \ln\left(\frac{1+x}{1-x}\right)$ for $-1 < x < 1$

Let $y = \tanh^{-1} x$

$$x = \tanh y$$

$$= \frac{e^y - e^{-y}}{e^y + e^{-y}}$$

$$x(e^y + e^{-y}) = e^y - e^{-y}$$

$$xe^y + xe^{-y} = e^y - e^{-y}$$

$$e^y (xe^y + xe^{-y}) = e^y (e^y - e^{-y})$$

$$xe^{2y} + x = e^{2y} - 1$$

$$0 = e^{2y} - xe^{2y} - x - 1$$

$$= e^{2y}(1-x) - x - 1$$

$$x+1 = e^{2y}(1-x)$$

$$e^{2y} = \frac{x+1}{1-x}$$

$$\ln(e^{2y}) = \ln\left(\frac{x+1}{1-x}\right)$$

$$2y = \ln\left(\frac{1+x}{1-x}\right)$$

$$y = \frac{1}{2} \ln\left(\frac{1+x}{1-x}\right)$$

Side Notes

- When doing proofs for inverse trig or inverse hyperbolic functions, start out with $y = (\text{inverse trig}) x$

$$x = \text{trig}(y)$$

- $\text{trig func}(\text{trig inverse func } x) = x$

Eg. $\sin(\sin^{-1} x) = x$

BUT! $\text{trig inverse func}(\text{trig func}) \neq x$

$$\sin^{-1}(\sin x) \neq x$$

- Principle functions:

$$\sin(x) \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$$

$$\cos(x) \in [0, \pi]$$

$$\tan(x) \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$

$$\csc(x) \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \quad x \neq 0$$

$$\sec(x) \in (0, \pi) \quad x \neq \frac{\pi}{2}$$

$$\cot(x) \in (0, \pi)$$

Summary of Derivatives

1. $\frac{d}{dx} c = 0$, c is a constant
2. $\frac{d}{dx} x = 1$
3. $\frac{d}{dx} cx = c$, c is a constant
4. $\frac{d}{dx} (f(x) \pm g(x)) = \frac{d}{dx} f(x) \pm \frac{d}{dx} g(x)$
5. $\frac{d}{dx} (f(x) \cdot g(x)) = f'(x)g(x) + f(x)g'(x)$
6. $\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$
7. $[(f(x))^n]' = n(f(x))^{n-1} \cdot f'(x)$
8. $(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$
9. If $f(x) = a^x$, for any $a > 0$, $a \neq 1$, then $f'(x) = a^x \cdot \ln a \cdot x'$
10. $(e^x)' = e^x \cdot x'$
11. If $f(x) = \log_b x$, for any $b > 0$, $b \neq 1$ then $f'(x) = \frac{1}{x \ln a} \cdot x'$
12. $f(x) = \ln(x) \rightarrow f'(x) = \frac{1}{x} \cdot x'$
13. $f(x) = \ln|x| \rightarrow f'(x) = \frac{x'}{x}$ * Special Case
14. $(\sin x)' = \cos x \cdot x'$
15. $(\cos x)' = -\sin x \cdot x'$
16. $(\tan x)' = \sec^2 x \cdot x'$
17. $(\sec x)' = \sec x \cdot \tan x \cdot x'$
18. $(\csc x)' = -\csc x \cdot \cot x \cdot x'$
19. $(\cot x)' = -\csc^2 x \cdot x'$
20. $(\sin^{-1} x)' = \frac{1}{\sqrt{1-x^2}} \cdot x'$
21. $(\cos^{-1} x)' = \frac{-1}{\sqrt{1-x^2}} \cdot x'$
22. $(\tan^{-1} x)' = \frac{1}{1+x^2} \cdot x'$
23. $(\sec^{-1} x)' = \frac{1}{|x| \sqrt{x^2-1}} \cdot x'$
24. $(\csc^{-1} x)' = \frac{-1}{|x| \sqrt{x^2-1}} \cdot x'$
25. $(\cot^{-1} x)' = \frac{-1}{1+x^2} \cdot x'$
- $(\sinh(x))' = \cosh(x) \cdot x'$
- $(\cosh(x))' = \sinh(x) \cdot x'$
- $(\tanh x)' = \text{sech}^2 x \cdot x'$
- $(\text{sech } x)' = -\text{sech } x \cdot \tanh x \cdot x'$
- $(\text{csch } x)' = -\text{csch } x \cdot \coth x \cdot x'$
- $(\coth x)' = -\text{csch}^2 x \cdot x'$
- $(\sinh^{-1} x)' = \frac{1}{\sqrt{1+x^2}} \cdot x'$
- $(\cosh^{-1} x)' = \frac{1}{\sqrt{x^2-1}} \cdot x'$
- $(\tanh^{-1} x)' = \frac{1}{1-x^2} \cdot x'$
- $(\text{sech}^{-1} x)' = \frac{-1}{x \sqrt{1-x^2}} \cdot x'$
- $(\text{csch}^{-1} x)' = \frac{-1}{|x| \sqrt{1+x^2}} \cdot x'$
- $(\coth^{-1} x)' = \frac{1}{1-x^2} \cdot x'$