

B52 Week 2 Notes

1. - Combinations:

The number of ways to choose k objects from a total of n objects is $\frac{n!}{(k)!(n-k)!}$ or nCk or $\binom{n}{k}$.

- Multinomial Coefficient:

The number of ways of dividing a group of n objects into k groups of sizes $n_1, n_2, \dots, n_k$ is $\frac{n!}{(n_1)!(n_2)! \dots (n_k)!}$ or $\binom{n}{n_1, n_2, \dots, n_k}$

- Ex. 1 One of my classes has 60 students. I want to divide my class into 2 groups, of even size, at random. If John and Jane are 2 students in my class, what is the prob that they will be in the same group?

Solution:

G_1 or G_2
 (J, J) (J, J)

Since the prob that Jack and Jane will both be in G_1 is $\frac{\binom{58}{28}}{\binom{60}{30}}$, the prob that they will be in G_2 together is also $\frac{\binom{58}{28}}{\binom{60}{30}}$.

$\therefore$ The answer is $\frac{\binom{58}{28}}{\binom{60}{30}} \times 2$

Conditional Probability:

- Let A and B be 2 events in a sample space, S . Then, the cond prob of A given B is defined by $P(A|B) = \frac{P(AB)}{P(B)}$, $P(B) > 0$.

$P(A|B)$ is read as "Probability of A given B ."

- E.g. 2 If A and B are events s.t. $P(A) = 0.5$, $P(B) = 0.3$ and $P(AB) = 0.1$, find:

a) $P(A|B)$

Solution:

$$\begin{aligned} P(A|B) &= \frac{P(AB)}{P(B)} \\ &= \frac{0.1}{0.3} \\ &= \frac{1}{3} \end{aligned}$$

b) $P(B|A)$

Solution:

$$\begin{aligned} P(B|A) &= \frac{P(AB)}{P(A)} \\ &= \frac{0.1}{0.5} \\ &= \frac{1}{5} \end{aligned}$$

c) $P(A|A \cup B)$

Solution:

$$\begin{aligned} P(A|A \cup B) &= \frac{P(A \cap (A \cup B))}{P(A \cup B)} \\ &= \frac{P(A)}{P(A) + P(B) - P(AB)} \\ &= \frac{0.5}{0.5 + 0.3 - 0.1} \\ &= \frac{0.5}{0.7} \\ &= \frac{5}{7} \end{aligned}$$

d) $P(A|AB)$

Solution:

$$\begin{aligned} P(A|AB) &= \frac{P(A \cap AB)}{P(AB)} \\ &= \frac{P(AB)}{P(AB)} \\ &= 1 \end{aligned}$$

Since AB is a subset of A, $A \cap AB = AB$.

- Ex. 3 A coin is flipped twice. $S = \{HH, HT, TH, TT\}$. What is the cond prob that both flips result in heads given that the first flip does?

Solution:

$$A = \{HH\}$$

$$B = \{HH, HT\}$$

$$P(A|B) = \frac{P(AB)}{P(B)}$$

$$= \frac{\frac{1}{4}}{\frac{2}{4}}$$

$$= \frac{1}{2}$$

- Thm of Total Prob:

Let $A_1, A_2, \dots, A_n$ be a partition of S and let B be any event in S .

Then, $P(B) = P(B|A_1)P(A_1) + P(B|A_2)P(A_2) + \dots + P(B|A_n)P(A_n)$.

Proof:

$$P(B) = P(B \cap A_1) + P(B \cap A_2) + \dots + P(B \cap A_n)$$

$$\text{However, } P(B|A_1) = \frac{P(B \cap A_1)}{P(A_1)}$$

$$\text{So, } P(B \cap A_1) = P(B|A_1) \cdot P(A_1)$$

$$\therefore P(B) = P(B|A_1) \cdot P(A_1) + P(B|A_2) \cdot P(A_2) + \dots + P(B|A_n) \cdot P(A_n)$$

QED

- Baye's Theorem:

Let A and B be any 2 events in S , then $P(A|B) = \frac{P(A) \cdot P(B|A)}{P(B)}$

Proof:

$$P(A|B) = \frac{P(AB)}{P(B)}$$

$$= \frac{P(B|A) \cdot P(A)}{P(B)} \quad \text{QED}$$

e) $P(A|A \cup B)$

Solution:

$$\begin{aligned} P(A|A \cup B) &= \frac{P((A|B) \cap (A \cup B))}{P(A \cup B)} \\ &= \frac{P(A|B)}{P(A \cup B)} \\ &= \frac{1}{7} \end{aligned}$$

- Thm: Conditional probability is a probability measure.

1. $0 \leq P(A|B) \leq 1$

2. $P(S|A) = 1$

3. $P(A_1 \cup A_2 \dots A_n | B) = P(A_1|B) + P(A_2|B) \dots + P(A_n|B)$

Proof of 1:

$$AB \subseteq B$$

$$P(AB) \leq P(B)$$

$$\frac{P(AB)}{P(B)} \leq \frac{P(B)}{P(B)}$$

$$\frac{P(AB)}{P(B)} \leq 1$$

$$P(B)$$

$$\therefore 0 \leq P(A|B) \leq 1$$

QED

Proof of 2:

$$P(S|A) = \frac{P(S \cap A)}{P(A)}$$

$$= \frac{P(A)}{P(A)}$$

$$= 1$$

$$= P(S)$$

QED

Proof of 3:

Suppose that $A_1, A_2, \dots, A_n$ are disjoint events.

$$\text{Then, } P(A_1 \cup A_2 \dots A_n) =$$

$$P(A_1) + P(A_2) + \dots + P(A_n).$$

$$P(A_1 \cup A_2 \dots A_n | B)$$

$$= \frac{P((A_1 \cup A_2 \dots A_n) \cap B)}{P(B)}$$

$$P(B)$$

$$= \frac{P(A_1 \cap B \cup A_2 \cap B \dots A_n \cap B)}{P(B)}$$

$$P(B)$$

$$= \frac{P(A_1 \cap B) + P(A_2 \cap B) + \dots + P(A_n \cap B)}{P(B)}$$

$$P(B)$$

$$= \frac{P(A_1 \cap B)}{P(B)} + \dots + \frac{P(A_n \cap B)}{P(B)}$$

$$= P(A_1|B) + P(A_2|B) + \dots + P(A_n|B)$$

QED

- E.g. 4 A population of voters consist of 40% Republicans and 60% Democrats. It is reported that 30% of R and 50% of D favour an election issue. A voter chosen at random favours the issue. What is the probability that this voter is a D?

Solution:

$$P(D) = \frac{60}{100}$$

$$= 0.6$$

$$P(R) = \frac{40}{100}$$

$$= 0.4$$

$$P(F|D) = \frac{50}{100}$$

$$= 0.5$$

$$P(F|R) = \frac{30}{100}$$

$$= 0.3$$

$$P(D|F) = \frac{P(D) \times P(F|D)}{P(F)}$$

$$= \frac{(0.6)(0.5)}{(0.42)}$$

$$= 0.714$$

$$P(F) = P(F|R) \cdot P(R) + P(F|D) \cdot P(D)$$

$$= (0.3)(0.4) + (0.5)(0.6)$$

$$= 0.42$$

3. Independent Events:

- Two events, A and B, in S are indep if $P(AB) = P(A) \times P(B)$.

$A \perp B$ denotes that A and B are indep.

$$- P(A|B) = \frac{P(AB)}{P(B)} \quad \text{If } P(B) > 0, \text{ then } A \perp B \text{ iff } P(A|B) = P(A).$$

$$= \frac{P(A) \times P(B)}{P(B)} \leftarrow \text{Since, if } A \perp B, \text{ then } P(AB) = P(A) \times P(B)$$

$$= P(A)$$

- Ex. 5 Suppose we roll a fair, six-sided die.

$$A = \{1, 3, 5\}$$

$$B = \{2, 4, 6\}$$

$$C = \{1, 2\}$$

$$D = \{1, 2, 3, 4, 5, 6\}$$

a) Are A and B indep?

Solution:

$$P(AB) = 0$$

$$\neq P(A) \times P(B)$$

$\therefore$ Not indep

b) Are A and C indep?

Solution:

$$P(A) = \frac{3}{6} = \frac{1}{2}$$

$$P(C) = \frac{2}{6} = \frac{1}{3}$$

$$P(AC) = \frac{1}{6}$$

$$= \frac{1}{2} \times \frac{1}{3}$$

$$= P(A) \times P(C)$$

$\therefore A \perp C$

- Note: If A and B are any 2 elements in S s.t. $P(A) > 0$ and $P(B) > 0$, if A and B are disjoint, then A and B are not indep.

Proof:

If A and B are disjoint,

$$P(AB) = 0$$

$$\neq P(A) \times P(B)$$

$\therefore$ A and B are not indep.

QED

- Thm: $A \perp B \rightarrow A \perp B^c$

Proof:

Suppose A and B are indep.

That means, $P(AB) = P(A) \cdot P(B)$

$$P(A \cap B^c) = P(A \setminus B)$$

$$= P(A) - P(AB)$$


$$= P(A) - P(A) \cdot P(B)$$

$$= P(A) [1 - P(B)]$$

$$= P(A) \cdot P(B^c), \text{ as wanted}$$

$$\text{i.e. } A \perp B^c$$

QED



The shaded region is $P(A \cap B^c)$.

- Thm: $A \perp B \rightarrow A^c \perp B^c$

Proof:

Suppose A and B are indep.

That means $P(AB) = P(A) \cdot P(B)$

$$P(A^c \cap B^c) = P(S) - P(A \cup B)$$

$$= 1 - P(A \cup B)$$

$$= 1 - [P(A) + P(B) - P(AB)]$$

$$= 1 - P(A) - P(B) + P(AB)$$

$$= (1 - P(A)) - P(B) + P(A) \cdot P(B)$$

$$= P(A^c) - P(B) + P(A) \cdot P(B)$$

$$= -[-P(A^c) + P(B) - P(A) \cdot P(B)]$$

$$= -[-P(A^c) + P(B) [1 - P(A)]]$$


$$= -[-P(A^c) + P(B) \cdot P(A^c)]$$

$$= P(A^c) - P(B) \cdot P(A^c)$$

$$= P(A^c) [1 - P(B)]$$

$$= P(A^c) \cdot P(B^c), \text{ as wanted}$$

QED



The shaded area is $A^c \cap B^c$ because $A^c \cap B^c$ is the area not in A and not in B .

- The events $A_1, A_2, A_3, \dots, A_n$ in S are indep if $P(A_{i1} \cap A_{i2} \cap \dots \cap A_{ik}) = P(A_{i1}) \times P(A_{i2}) \times \dots \times P(A_{ik})$ for each finite subcollection of $A_{i1}, A_{i2}, \dots, A_{ik}$.

- E.g. 6 Suppose we roll a 4-sided die.
 $S = \{1, 2, 3, 4\}$
 $A = \{1, 2\}$
 $B = \{1, 3\}$
 $C = \{1, 4\}$

a) Are A and B indep?

Solution:

$$P(A) = \frac{1}{2}$$

$$P(B) = \frac{1}{2}$$

$$P(AB) = \frac{1}{4}$$

$$= \frac{1}{2} \times \frac{1}{2}$$

$$= P(A) \cdot P(B)$$

$\therefore$ Yes, $A \perp B$

b) Are B and C indep?

Solution:

$$P(B) = \frac{1}{2}$$

$$P(C) = \frac{1}{2}$$

$$P(BC) = \frac{1}{4}$$

$$= \frac{1}{2} \times \frac{1}{2}$$

$$= P(B) \cdot P(C)$$

$\therefore$ Yes, $B \perp C$

c) Are A and C indep?

Solution:

$$P(A) = \frac{1}{2}$$

$$P(C) = \frac{1}{2}$$

$$P(AC) = \frac{1}{4}$$

$$= \frac{1}{2} \times \frac{1}{2}$$

$$= P(A) \cdot P(C)$$

$\therefore$ Yes, $A \perp C$

d) Are A and B and C indep?

Solution:

$$P(A) = P(B) = P(C) = \frac{1}{2}$$

$$P(A \cap B \cap C) = \frac{1}{4} \quad (\text{Because } P(A \cap B \cap C) = \{1\})$$

$$\neq \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2}$$

$\therefore$ A and B and C are not indep.

Note: Pair-wise indep does not imply indep.