

Master Theorem

Introduction:

- Master's Thm states:

Let $a \geq 1$ and $b > 1$ be constants.

Let $f(n)$ be an asymptotically positive function.

Let $T(n) \leq aT\left(\frac{n}{b}\right) + O(f(n))$ be a recurrence relation.

Then:

1. If $f(n) = O(n^{\log_b a - \epsilon})$ for some constant $\epsilon > 0$, then $T(n) = O(n^{\log_b a})$

2. If $f(n) = \Theta(n^{\log_b a} \cdot \log^k n)$ for some constant $k \geq 0$, then $T(n) = O(n^{\log_b a} \log^{k+1} n)$

3. If $f(n) = \Omega(n^{\log_b a + \epsilon})$ for some constant $\epsilon > 0$ and f satisfies the **regularity condition** then $T(n) = O(f(n))$

The **regularity condition** states that:

For some constant $c < 1$ and all sufficiently large n , $a \cdot f\left(\frac{n}{b}\right) \leq c \cdot f(n)$

Examples:

1. $T(n) \leq 9T\left(\frac{n}{3}\right) + n$

$$a=9, b=3, f(n)=n, \log_b a = \log_3 9 = 2$$

$$\text{We see that } n^{\log_b a} = n^2.$$

Hence, $f(n) = O(n^{\log_b a - \epsilon})$, where $\epsilon = 1$ in this case.

$$\therefore T(n) = O(n^2)$$

2

$$2. T(n) \leq T\left(\frac{2n}{3}\right) + 1$$

$$a=1, b=\frac{2}{3}, f(n)=1, \log_b a = 0 \\ n^{\log_b a} = n^0 = 1$$

We see that $f(n) = n^{\log_b a}$. We can see that case 2 applies if $k=0$.

$$\therefore T(n) = O(n \log n)$$

$$3. T(n) \leq 2T\left(\frac{n}{2}\right) + n \lg(n)$$

$$a=2, b=2, f(n)=n \lg(n), \log_b a = \log_2 2 = 1$$

None of the cases apply here.

While $f(n)$ is larger than $n^{\log_b a}$ or n , it is not polynomially larger.

$$\frac{n \lg n}{n} = \lg(n)$$

$$\text{For case 3, we need } f(n) = \Omega(n^{\log_b a + \epsilon}) \\ = \Omega(n^{\log_b a} \cdot n^\epsilon)$$

and $\lg(n)$ is asymptotically less than n^ϵ for any positive ϵ .

$$4. T(n) \leq 2T\left(\frac{n}{2}\right) + O(n)$$

Note: This is the run time for the divide and conquer soln to max subarray.

$$a=2, b=2, f(n)=O(n), \log_b a = 1$$

This would fall under case 2, for $k=0$.

$$\therefore T(n) = O(n \lg n)$$

$$5. T(n) \leq 7(T(\frac{n}{2})) + O(n^2)$$

Note: This is the run time for Strassen's algo.

$$a=7, b=2, f(n)=O(n^2), \log_b a = \log_2 7 \\ \approx 2.8$$

The first case applies as $f(n) = O(n^{\log_2 7 - \epsilon})$
where $\epsilon = 0.8$.

$$\therefore T(n) = O(n^{\log_b a}) \\ = O(n^{\log_2 7})$$